

**G006
Anilines**

Results of Testing

Chemical Name	CAS No.	Study Code/Type	Protocol/Guideline	Species	Exposure	Dose/Concentration	No. per Group	Results	Reference
4-Nitroaniline	100-01-6	HEGTOXCHRM Mammalian bone marrow micronucleus assay	Non-TSCA Protocol/Guideline (docket OPTS-42054B)	mice	intraperitoneal injection, 2x, 24 hours apart	0, 80, 400, 800 mg/kg/day	5 to 6/sex	No evidence of clastogenicity was found in any treatment group.	54 FR 42034; 10/13/89 OTS0532109
4-Chloroaniline	106-47-8	HEGTOXCHRM Mammalian bone marrow micronucleus assay	Non-TSCA Protocol/Guideline (docket OPTS-42054B)	mice	oral (gavage), single dose	0, 50, 100, 200 mg/kg body weight	5 male; 5 female	The incidence of micronucleated polychromatic erythrocytes in the test animals treated with 4-chloroaniline were within normal range. The number of normochromatic erythrocytes containing micronuclei was not increased. The ratio of polychromatic/normochromatic erythrocytes in both male and female test animals remained unaffected. Results indicated that the test material was not mutagenic.	53 FR 45385; 11/9/88 OTS0519119
Aniline	62-53-3	EEATOX Acute aquatic invertebrate toxicity	Non-TSCA Protocol/Guideline (docket OPTS-42054B)	<i>Gammarus fasciatus</i> (amphipod)	flow-through, 96 hours	0.18, 0.38, 0.70, 1.4, 2.7 mg/L (measured)	20 (10/replicate)	Exposure of the test animals to the test material (aniline) resulted in a 96-hour LC ₅₀ of 2.3 mg/L (1.9 to 3.1 mg/L). The no-observed-effect concentration (NOEC) based on survival was 1.4 mg/L.	54 FR 25167; 6/13/89 OTS0519116
Aniline	62-53-3	EECTOX Chronic aquatic toxicity - crustacean	Non-TSCA Protocol/Guideline (docket OPTS-42054B)	<i>Daphnia magna</i>	flow-through, 21 days	0.006-0.040 (measured)	20 (10/replicate)	No effects were noted at 0.016 mg/L. At 0.027 mg/L and higher, reproduction was significantly decreased as compared to controls. The maximum allowable toxicant concentration (MATC) was 0.021 mg/L.	54 FR 33772; 8/16/89 OTS0532105
Aniline	62-53-3	EECTOX Chronic aquatic toxicity - crustacean	Non-TSCA Protocol/Guideline (docket OPTS-42054B)	<i>Daphnia magna</i>	flow-through, 21 days	0.006-0.040 mg/L	20 (10/replicate)	Decreased reproduction occurred at 0.027 mg/L and higher. No effects were noted at 0.016 mg/L. The MATC was 0.021 mg/L, measured concentration.	54 FR 33773; 8/16/89 OTS0532105
Aniline	62-53-3	HEGTOXCHRM Mammalian bone marrow micronucleus assay	Non-TSCA Protocol/Guideline (docket OPTS-42054B)	mice	intraperitoneal injection, 2x, 24 hours apart	0, 30, 100, 300 mg/kg/day	3/sex	Increased incidence of micronucleated polychromatic erythrocytes were seen in the high dose groups for both sexes.	54 FR 33773; 8/16/89 OTS0532103
2-Nitroaniline	88-74-4	HEGTOXCHRM Mammalian bone marrow micronucleus assay	Non-TSCA Protocol/Guideline (docket OPTS-42054B)	mice	intraperitoneal injection, 2x, 24 hours apart	0, 50, 250, 500 mg/kg/day	5 to 6/sex	No evidence of clastogenicity was noted in any dose group.	54 FR 42034; 10/13/89 OTS0532108
2-Chloroaniline	95-51-2	EEATOX Acute fish toxicity	Non-TSCA Protocol/Guideline (docket OPTS-42054B)	Rainbow trout	flow-through, 96 hours	0.30, 0.58, 1.1, 2.0, 4.3 mg/L (measured)	20 (10/replicate)	The test material had an LC ₅₀ value (and a 95% confidence limit) of 1.0 mg/L (0.82 to 1.4 mg/L). Altered body coloration and erratic swimming were noted.	54 FR 25167; 6/13/89 OTS0519118

G006 Anilines

Chemical Name	CAS No.	Study Code/Type	Protocol/Guideline	Species	Exposure	Dose/Concentration	No. per Group	Results	Reference
2-Chloroaniline	95-51-2	EEATOX Acute invertebrate toxicity	Non-TSCA Protocol/Guideline (docket OPTS-42054B)	<i>Gammarus fasciatus</i> (amphipod)	flow-through, 96 hours	0, 0.72, 1.2, 2.0, 3.8, 8.2 mg/L	Not specified	Exposure to the test material resulted in a 96-hour LC ₅₀ value of 5.4 mg/L (2.9 to 0.62 mg/L). The no-observed-effect concentration based on survival was 3.8 mg/L. Test animals exposed to 8.2 mg/L exhibited lethargy and immobilization.	54 FR 25167; 6/13/89 OTS0519118
2-Chloroaniline	95-51-2	EECLIF Fish early life stage test	Non-TSCA Protocol/Guideline (docket OPTS-42054B)	rainbow trout	flow-through, 105 days	0.0037-1.4 mg/L	60/concentration (30/replicate)	The NOEC was 0.0037 mg/L, and the lowest effect concentration was 0.012 mg/L (growth in length). The MATC was 0.0067 mg/L.	54 FR 33772; 8/16/89 OTS0532104
2-Chloroaniline	95-51-2	EECTOX Chronic aquatic toxicity - crustacean	Non-TSCA Protocol/Guideline (docket OPTS-42054B)	<i>Daphnia magna</i>	static, 21 days	0.013-0.19 mg/L	Not specified	Survival was decreased at 0.19 mg/L, and at 0.046 mg/L and higher, decreased total young produced and offspring per surviving adult was noted.	54 FR 33773; 8/16/89 OTS0532104
2-Chloroaniline	95-51-2	EECTOX Chronic aquatic toxicity - crustacean	Non-TSCA Protocol/Guideline (docket OPTS-42054B)	<i>Daphnia magna</i>	flow-through, 21 days	0.013-0.19 mg/L (measured)	10 (10/replicate)	The no-effect level was 0.025 mg/L. At 0.19 mg/L, survival was significantly decreased over controls, and at 0.046 mg/L and higher reproduction was decreased. The MATC was 0.025 mg/L.	54 FR 33772; 8/16/89 OTS0532104
2-Chloroaniline	95-51-2	HEGTOXCHRM Mammalian bone marrow micronucleus assay	Non-TSCA Protocol/Guideline (docket OPTS-42054B)	mice	intraperitoneal injection, 2x, 24 hours apart	0, 20, 70, 200 mg/kg/day	4 or 5/sex	No evidence of increased micronucleated polychromatic erythrocytes were seen at any test level.	54 FR 39806; 9/28/89 OTS0532107
3,4-Dichloroaniline	95-76-1	HEGTOXCHRM Mammalian bone marrow micronucleus assay	Non-TSCA Protocol/Guideline (docket OPTS-42054B)	mice	intraperitoneal injection, 2x, 24 hours apart	0, 20, 70, 200 mg/kg/day	5/sex	No evidence of clastogenicity was found in any treatment group.	54 FR 43482; 10/25/89 OTS0532110
2,4-Dinitroaniline	97-02-9	HEGTOXCHRM Mammalian bone marrow micronucleus assay	Non-TSCA Protocol/Guideline (docket OPTS-42054B)	mice	oral (gavage), single dose	0, 37.5, 75, 150 mg/kg body weight	5/sex	Test animals treated with 2,4-dinitroaniline had an incidence of micronucleated polychromatic erythrocytes within normal range. The ratio of polychromatic normochromatic erythrocytes in both male and female test animals remained unaffected. Results indicated that the test material was not mutagenic.	53 FR 45385; 11/9/88 OTS0519120
2,6-Dichloro-4-nitroaniline	99-30-9	EEATOX Algae acute toxicity	Non-TSCA Protocol/Guideline (docket OPTS-42054B)	<i>Selenastrum capricornutum</i> (algae)	static, 96 hours	0.9, 1.3, 1.9, 2.8, 4.2 mg/L (measured)	Not applicable	Exposure to the test material (2,6-dichloro-4-nitroaniline) resulted in a 96-hour EC ₅₀ value of 2.6 mg/L. The no-observed-effect concentration was 0.9 mg/L.	54 FR 25167; 6/13/89 OTS0519117
2,6-Dichloro-4-nitroaniline	99-30-9	EEATOX Acute invertebrate toxicity	Non-TSCA Protocol/Guideline (docket OPTS-42054B)	<i>Daphnia magna</i>	flow-through, 48 hours	0.86-5.0 mg/L (nominal)	20 (10/replicate)	Exposure to 2,6-dichloro-4-nitroaniline produced a 48-hour EC ₅₀ value greater than 4.4 mg/L (the highest concentration due to solubility limitations). No evidence of acute toxicity was seen at any test concentration.	54 FR 25167; 6/13/89 OTS0519117

**G006
Anilines**

Chemical Name	CAS No.	Study Code/Type	Protocol/Guideline	Species	Exposure	Dose/Concentration	No. per Group	Results	Reference
2,6-Dichloro-4-nitroaniline	99-30-9	EECLIF Fish early life stage test	Non-TSCA Protocol/Guideline (docket OPTS-4453B)	rainbow trout	flow-through, 91 days	0.011-0.19 mg/L	60/ concentration	Decreased larval survival was noted at 0.024 mg/L and higher. The NOEC was 0.011 mg/L and the MATC was 0.016 mg/L.	54 FR 30605; 8/31/89, Docket OPTS-44536